




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

Predictors of silent hypoxia in hospitalized patients with COVID-19 in Japan

Kayoko Hayakawa, Shinichiro Morioka, Yusuke Asai, Shinya Tsuzuki, Gen Yamada, Setsuko Suzuki, Nobuaki Matsunaga, Norio Ohmagari



PII: S1341-321X(22)00176-3

DOI: <https://doi.org/10.1016/j.jiac.2022.06.001>

Reference: JIC 1934

To appear in: *Journal of Infection and Chemotherapy*

Received Date: 3 April 2022

Revised Date: 19 May 2022

Accepted Date: 2 June 2022

Please cite this article as: Hayakawa K, Morioka S, Asai Y, Tsuzuki S, Yamada G, Suzuki S, Matsunaga N, Ohmagari N, Predictors of silent hypoxia in hospitalized patients with COVID-19 in Japan, *Journal of Infection and Chemotherapy* (2022), doi: <https://doi.org/10.1016/j.jiac.2022.06.001>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Japanese Society of Chemotherapy and The Japanese Association for Infectious Diseases.
Published by Elsevier Ltd. All rights reserved.

Note

Title: Predictors of silent hypoxia in hospitalized patients with COVID-19 in Japan

Kayoko Hayakawa^{a,b}, Shinichiro Morioka^{a,b,c}, Yusuke Asai^b, Shinya Tsuzuki^{a,b,d}, Gen Yamada^a, Setsuko Suzuki^a, Nobuaki Matsunaga^b, and Norio Ohmagari^{a,b}

^aDisease Control and Prevention Center, National Center for Global Health and Medicine, Tokyo, Japan

^bAMR Clinical Reference Center, National Center for Global Health and Medicine, Tokyo, Japan

^cEmerging and Reemerging Infectious Diseases, Graduate School of Medicine, Tohoku University, Sendai, Japan

^dFaculty of Medicine and Health Sciences, University of Antwerp, Antwerp, Belgium.

Running title: Silent hypoxia in COVID-19 patients

Corresponding author:

Kayoko Hayakawa

Address: 1-21-1 Toyama, Shinjuku-ku, Tokyo 162-8655, Japan

Telephone: +81-3-3202-7181

Fax: +81-3-6228-0738

E-mail: khayakawa@hosp.ncgm.go.jp

Abstract

Introduction

Silent hypoxia (SH) is common in patients with coronavirus disease (COVID-19) in Japan and other countries. Early identification of SH is important as more treatment options for COVID-19 have become available. This study aimed to identify predictors of SH using a nationwide COVID-19 registry of hospitalized patients.

Methods

Adult patients who were admitted to hospital with COVID-19 between January 2020 and June 2021 and who were hypoxic on admission (SpO_2 : 70–94%), not transferred from another facility, and who did not have disturbance of consciousness, confusion, or dementia, were included. SH was defined as hypoxia in the absence of shortness of breath/dyspnea upon admission. Predictors of SH were identified using univariable and multivariable logistic regression.

Results

The study included 1,904 patients, of whom 990 (52%) satisfied the criteria for SH. Compared to patients without SH, patients with SH were older, more likely to be female, and had a slightly higher SpO_2 on admission. Compared to patients without SH, patients with SH had a lower prevalence of chronic lung disease (CLD) other than chronic obstructive pulmonary disease (COPD), asthma, and obesity. Multivariable analysis revealed that the independent predictors of SH were older age, a shorter interval from symptom onset to admission, higher SpO_2 , and an absence of CLD or COPD.

Conclusions

The absence of underlying lung disease and older age were important predictors of SH. The results of this study, which is the largest such study reported to date in Japan, may help clarify the mechanism of SH.

Keywords: silent hypoxia, COVID-19, infection

Silent hypoxia (SH), also known as happy hypoxia, has been frequently observed in COVID-19 patients Japan and other countries [1-4]. The early identification of SH and initiation of therapeutic interventions are important given that more treatment options for coronavirus disease-2019 (COVID-19) have now become available. This study was conducted to identify the predictive factors for SH using the nationwide COVID-19 registry of hospitalized patients (COVIREGI-JP).

The patients who fulfilled the following criteria were included in the analysis: (1) ≥ 20 years of age, (2) no oxygen usage on admission, (3) $70\% < \text{SpO}_2, < 94\%$ on room air, (4) no disturbance of consciousness (AVPU scale: alert or verbal), (5) no confusion or dementia, and (6) not a transferred patient. SH was defined as the absence of shortness of breath (SOB)/dyspnea upon admission. The data of patients, admitted between January 2020 and June 2021, which were fixed by September 30, 2021, were used as previously described [5]. All analyses were performed using IBM SPSS 25. Univariable and multivariable analyses were performed using logistic regression, and independent predictors for the silent hypoxia were identified. Two-sided P value of $< .05$ was considered statistically significant. This study was approved by the Ethics Committee of the National Center for Global Health and Medicine (NCGM-G-004147-00).

The study included 1,904 patients, of whom 990 (52%) satisfied the criteria for SH. In univariable analysis, compared to the non-SH group, the SH group had more female and

older adult patients (Table). Alcohol consumption was more prevalent in the non-SH group than in the SH group. The patients from the SH group had a slightly higher SpO₂ on admission, as well as a slightly lower temperature and heart rate than those in the non-SH group. The prevalence of chronic lung disease (CLD) other than chronic obstructive pulmonary disease (COPD), asthma, and obesity, was lower in the SH group than in the non-SH group. The days from symptom onset (DSO) were shorter in the SH group than in the non-SH group. Multivariate analysis revealed that the independent predictors of SH were older age, shorter DSO, higher SpO₂, and not having CLD or COPD.

The results were partially similar to those in the study by García-Grimshaw *et al.*, which identified DSO as a predictor of SH [6]. However, they were not concordant with the report by Alhusain *et al.*, which did not include DSO or vitals on admission [7]. Both studies used different definitions of hypoxia and included symptoms as predictors. In the present study, although symptoms were excluded to avoid confounding effects, more comorbidities were considered. The correlation between the risk factors for severe COVID-19 and the predictors of SH was minimal [8].

Based on our results, patients with COPD and CLD were more likely to complain of SOB. Oxygen-requiring patients on admission were not included in the study; therefore, patients with advanced COPD or CLD were likely excluded. These findings suggest that patients with underlying pulmonary diseases that are not sufficiently advanced for them to be

accustomed to hypoxia, are less likely to develop SH because they tend to be more aware of their respiratory status.

Various hypotheses regarding the pathomechanism of SH have been proposed [9-11]. The lung perfusion, sensory feedback, and central neural regulation of breathing are likely to be affected in patients with underlying lung abnormalities. The findings of the present study require further basic investigation and validation in non-Japanese cohorts.

Limitations of this study include the use of registry data, which may have resulted in selection bias, as previously reported [5]. Although we performed multivariable analysis, there may be some residual confounding.

In conclusion, in a large cohort of patients hospitalized with COVID-19, the absence of underlying lung disease and age were important predictors of SH. The results of this study, which included the largest number of reported cases, may help clarify the mechanism of SH.

Potential conflicts of interest: The authors declare no conflicts of interest. All authors have submitted the ICMJE form for the disclosure of potential conflicts of interest. Conflicts that the editors considered relevant to the content of the manuscript have been disclosed.

Funding: This research was funded by the Health and Labor Sciences Research grant, “Research for risk assessment and implementation of crisis management functions for

emerging and re-emerging infectious diseases.”

K.H was the chief investigator and responsible for the data analysis. S.M contributed to the study design and ethical approval. Y.A, S.T, and G.Y reviewed the statistical analyses. K.H drafted the manuscript. All authors contributed to the reviewing and finalization of the manuscript.

Acknowledgments. The authors thank all participating facilities for their care of patients with COVID-19 and their cooperation in data entry into the registry.

References

1. Akiyama Y, Morioka S, Asai Y, Sato L, Suzuki S, Saito S, Matsunaga N, Hayakawa K and Ohmagari N. Risk factors associated with asymptomatic hypoxemia among covid-19 patients: A retrospective study using the nationwide japanese registry, COVIREGI-JP. *J Infect Public Health*. 2022;15:312-14.
2. Busana M, Gasperetti A, Giosa L, Forleo GB, Schiavone M, Mitacchione G, Bonino C, Villa P, Galli M, Tondo C, Saguner A, Steiger P, Curnis A, Dello Russo A, Pugliese F, Mancone M, Marini JJ and Gattinoni L. Prevalence and outcome of silent hypoxemia in covid-19. *Minerva Anesthesiol*. 2021;87:325-33.
3. Okuhama A, Ishikane M, Hotta M, Sato L, Akiyama Y, Morioka S, Suzuki S, Tajima T, Yamamoto M, Teruya K, Izumi S and Ohmagari N. Clinical and radiological findings of silent hypoxia among covid-19 patients. *J Infect Chemother*. 2021;27:1536-38.
4. Ribeiro A, Mendonca M, Sabina Sousa C, Trigueiro Barbosa M and Morais-Almeida M. Prevalence, presentation and outcomes of silent hypoxemia in covid-19. *Clin Med Insights Circ Respir Pulm Med*. 2022;16:11795484221082761.
5. Matsunaga N, Hayakawa K, Terada M, Ohtsu H, Asai Y, Tsuzuki S, Suzuki S, Toyoda A, Suzuki K, Endo M, Fujii N, Suzuki M, Saito S, Uemura Y, Shibata T, Kondo M, Izumi K, Terada-Hirashima J, Mikami A, Sugiura W and Ohmagari N. Clinical epidemiology of hospitalized patients with covid-19 in Japan: Report of the COVID-19 registry japan. *Clin*

Infect Dis. 2021;73:e3677-e3689.

6. Garcia-Grimshaw M, Flores-Silva FD, Chiquete E, Cantu-Brito C, Michel-Chavez A, Vigueras-Hernandez AP, Dominguez-Moreno R, Chavez-Martinez OA, Sanchez-Torres S, Marche-Fernandez OA and Gonzalez-Duarte A. Characteristics and predictors for silent hypoxemia in a cohort of hospitalized covid-19 patients. *Auton Neurosci*. 2021;235:102855.
7. Alhusain F, Alromaih A, Alhajress G, Alsaghyir A, Alqobaisi A, Alaboodi T and Alsalamah M. Predictors and clinical outcomes of silent hypoxia in covid-19 patients, a single-center retrospective cohort study. *J Infect Public Health*. 2021;14:1595-99.
8. Terada M, Ohtsu H, Saito S, Hayakawa K, Tsuzuki S, Asai Y, Matsunaga N, Kutsuna S, Sugiura W and Ohmagari N. Risk factors for severity on admission and the disease progression during hospitalisation in a large cohort of patients with covid-19 in japan. *BMJ Open*. 2021;11:e047007.
9. Herrmann J, Mori V, Bates JHT and Suki B. Modeling lung perfusion abnormalities to explain early covid-19 hypoxemia. *Nat Commun*. 2020;11:4883.
10. Simonson TS, Baker TL, Banzett RB, Bishop T, Dempsey JA, Feldman JL, Guyenet PG, Hodson EJ, Mitchell GS, Moya EA, Nokes BT, Orr JE, Owens RL, Poulin M, Rawling JM, Schmickl CN, Watters JJ, Younes M and Malhotra A. Silent hypoxaemia in covid-19 patients. *J Physiol*. 2021;599:1057-65.
11. Dhont S, Derom E, Van Braeckel E, Depuydt P and Lambrecht BN. The pathophysiology

of 'happy' hypoxemia in covid-19. *Respir Res.* 2020;21:198.

12. Charlson ME, Pompei P, Ales KL and MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. *Journal of chronic diseases.* 1987;40:373-83.

Journal Pre-proof

Table. Predictors for silent hypoxia on admission among hypoxic COVID-19 patients

Parameters	Silent	Non-Silent	Univariable		Multivariable analysis	
	Hypoxia ^a	Hypoxia ^a	analysis			
	(n=990)	(n=914)	OR	P value ^b	OR	P value ^b
Demographics						
Age (years), median (IQR)	71 (60–80)	65 (54–76)	1.02 (1.02-1.03)	<0.001	1.02 (1.01-1.03)	0.002
Male sex	617 (62.3%)	663 (72.7%)	0.62 (0.51–0.75)	<0.001	0.8 (0.58–1.11)	0.179
Japanese race	950 (97.2%)	870 (96%)	1.46 (0.88–2.42)	0.147	1.5 (0.74–3.05)	0.258

Current or previous smoker	429 (51.5%)	434 (54.5%)	0.89 (0.73–1.08)	0.232	1.01 (0.76–1.35)	0.93
Alcoholic beverage drinker	382 (51.8%)	406 (57.5%)	0.8 (0.65–0.98)	0.031	1.02 (0.76–1.36)	0.92
Days from symptom onset, median (IQR)	6 (3–8)	6 (4–9)	0.93 (0.91–0.96)	<0.001	0.94 (0.91–0.98)	0.001
Vital signs on admission						
SpO ₂ , median (IQR)	92 (91–93)	91 (89–93)	1.16 (1.12–1.2)	<0.001	1.14 (1.09–1.19)	<0.001
Temperature in Celsius, median (IQR)	37.4 (36.8–38.1)	37.5 (36.9–38.3)	0.88 (0.8–0.97)	0.007	1.06 (0.92–1.22)	0.392

Respiratory rate, median (IQR)	20 (17–22)	21 (18–24)	1.00 (1.00-1.00)	0.512	1.00 (1.00–1.00)	0.551
Heart rate, median (IQR)	89 (80–101)	92 (82–103)	0.99 (0.98-0.99)	<0.001	1.00 (0.99–1.01)	0.472
Comorbidities^c						
Myocardial infarction	33 (3.3%)	32 (3.5%)	0.95 (0.58–1.56)	0.84	1.00 (0.46–2.14)	0.992
Congestive heart failure	41 (4.1%)	24 (2.6%)	1.6 (0.96–2.67)	0.071	1.78 (0.79–4.02)	0.168
Peripheral vascular disease	18 (1.8%)	23 (2.5%)	0.72 (0.39-1.34)	0.296	0.43 (0.15–1.26)	0.125

Cerebrovascular disease	86 (8.7%)	55 (6%)	1.49 (1.05–2.11)	0.027	1.14 (0.66–1.98)	0.64
Chronic lung disease (excluding COPD)	23 (2.3%)	41 (4.5%)	0.51 (0.3–0.85)	0.010	0.27 (0.11–0.66)	0.004
COPD	50 (5.1%)	62 (6.8%)	0.73 (0.5–1.07)	0.11	0.35 (0.18–0.68)	0.002
Asthma	42 (4.2%)	65 (7.1%)	0.58 (0.39–0.86)	0.007	0.67 (0.39–1.16)	0.154
Liver disease	33 (3.3%)	29 (3.2%)	1.05 (0.63–1.75)	0.844	1.08 (0.48–2.43)	0.85
Peptic ulcer disease	11 (1.1%)	7 (0.8%)	1.46	0.439	1.29	0.738

			(0.56–3.77)		(0.3–5.58)	
			0.9			
Diabetes mellitus	271 (27.4%)	269 (29.4%)	(0.74–1.1)	0.32	0.85 (0.63–1.16)	0.303
			0.6		0.98	
Obesity ^d	71 (7.2%)	105 (11.5%)	(0.43–0.82)	0.001	(0.64–1.51)	0.927
			1.76 (0.78–3.96)	0.174	3.46	
Severe renal dysfunction	17 (1.7%)	9 (1%)			(0.69–17.25)	0.13
			1		0.65	
Solid tumors	51 (5.2%)	47 (5.1%)	(0.67–1.51)	0.993	(0.33–1.27)	0.205
			1.05		0.75	
Metastatic solid tumors	17 (1.7%)	15 (1.6%)	(0.52–2.11)	0.897	(0.27–2.14)	0.594

Leukemias or lymphomas	6 (0.6%)	10 (1.1%)	0.55 (0.2–1.52)	0.251	0.36 (0.07–1.86)	0.221
Collagen disease	16 (1.6%)	20 (2.2%)	0.73 (0.38–1.43)	0.362	0.65 (0.23–1.84)	0.415
Hypertension	465 (47%)	401 (43.9%)	1.13 (0.95–1.36)	0.175	1.01 (0.76–1.35)	0.936
Dyslipidemia	222 (22.4%)	218 (23.9%)	0.92 (0.75–1.14)	0.461	1.05 (0.77–1.44)	0.758

^aPresented as number (%) unless otherwise indicated. ^bTwo-sided P value of < 0.05 was considered statistically significant (indicated as bold text). ^cDefinitions were based on their Charlson Comorbidity Index scores, unless otherwise specified [12]. ^dBased on the physician's diagnosis. Abbreviations. COPD, chronic obstructive pulmonary disease; IQR, interquartile range; OR, odds ratio.